

Spatial Coherence and Crest-Length Statistics of Waves in Deep Water

Steve Elgar
Woods Hole Oceanographic Institution
Applied Ocean Physics & Engineering, MS #11
Woods Hole, MA 02543
phone: (508) 289-3614 fax: (508) 457-2194 e-mail: elgar@whoi.edu

T.H.C. Herbers
Department of Oceanography, Code OC/He
Naval Postgraduate School
Monterey, CA 93943-5123
phone: (408) 656-2971 fax: (408) 656-2712 e-mail: herbers@oc.nps.navy.mil

Award Number: N00014-99-11070

LONG-TERM GOALS

The long-term objective is to determine how nonlinear interactions and directional spreading affect the spatial coherence and crest-length statistics of ocean surface gravity waves.

OBJECTIVES

To investigate the spatial coherence and statistical distribution of crest lengths of deep water ocean waves we have been developing and verifying with field observations the methodology to simulate two-dimensional sea surfaces with a specified frequency-directional spectrum and local nonlinearity described by theory. Our specific objectives are to

- simulate realizations of the two-dimensional sea surface given a specified frequency-directional spectrum and incorporating local nonlinearity based on second-order theory
- estimate crest-length statistics from the simulated sea surfaces for a range of wave conditions
- compare simulated with remotely-sensed sea surfaces

APPROACH

We have developed the methodology to simulate realizations of a nonlinear sea surface with a specified ('target') frequency-directional or wavenumber spectrum and a specified bispectrum. The bispectrum describes statistically the phase relationships between triads of nonlinearly interacting waves, for example the quadratic difference interaction that couples two swell waves (with wavenumbers k_p and $k_p + \delta$, where δ is a small number, and a low frequency long wave with wavenumber δ), or sum interactions that couple swell with higher harmonics. The simulation algorithm accounts for interactions between waves traveling in different directions. The target wavenumber spectrum can be obtained from observations, the output of a wave prediction model, or theory. The target bispectrum can be obtained from observations or from second-order nonlinear theory given the wavenumber spectrum.

DTIC QUALITY INSPECTED 4

20000911 116

To test second-order nonlinear theory, statistics of the observed wave field (eg, higher-order spectra, sea-surface elevation skewness, crest length) are compared with statistics from sea surfaces simulated from the observed wavenumber spectrum and the corresponding bispectrum consistent with second-order theory.

WORK COMPLETED

Software to estimate higher-order spectra of sea surfaces has been produced. In addition, the theory to determine the bispectrum from the wavenumber spectrum estimated with observations from a moving platform (eg, an airplane) has been developed. Simulations with narrow (Figure 1) and broad (Figure 2) frequency-directional spectra show differences in the corresponding sea surfaces. In particular, the groups are larger (more big waves in a row) and the crests are longer for narrow band sea surfaces than for the broad band waves.

Wavenumber and higher-order spectra have been estimated from LIDAR observations (provided by P. Hwang, NRL) of the sea surface. Theoretical bispectra have been calculated from the observed wavenumber spectra, and both theoretical and observed bispectra have been used as input to simulate realizations of the sea surface.

RESULTS

The bispectrum estimated from the observed wavenumber spectrum and second-order nonlinear theory compares well with the bispectrum estimated directly from LIDAR images of approximately 1 m high waves in 20-30 m water depth near the North Carolina coast. Third-order statistics of the sea surfaces obtained from numerical simulations using the observed wavenumber spectrum and either the observed or theoretical bispectrum compare well with the statistics of the observed sea surface, verifying both the simulation technique and second-order nonlinear theory.

IMPACT/APPLICATIONS

An impact of this research is the verification of second-order theory for directionally spread waves, allowing realistic nonlinear sea surfaces to be simulated given the wavenumber spectrum.

TRANSITIONS

RELATED PROJECTS

The research performed here is related to our investigations of the evolution of waves across the continental shelf (SHOWEX) and the nearshore and surfzone (SandyDuck).

PUBLICATIONS

Herbers, T.H.C., Steve Elgar, N.A. Sarap, and R.T. Guza, 2000. Dispersion properties of surface gravity waves in shallow water, *J. Physical Oceanography*, submitted.




Figure 1. Sea surfaces consisting of lowest-order wind generated waves with broad (upper) and narrow (lower) frequency-directional spectra, and second-order nonlinear waves consistent with theory. The amount of nonlinear coupling for each pair of wind waves and the corresponding sum or difference second-order waves is a function of the wavenumber spectrum and the frequencies of the first-order wind waves. The significant wave height is 5.7 m. The colors are sea-surface elevation (red is wave crests, dark blue is troughs). The small panels on the left and bottom of the color contours are 1D cuts through the center of the sea surface (dotted lines on the sea surfaces show the locations of the 1D surfaces).

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (07-04-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 2 September 2000	3. REPORT TYPE AND DATES COVERED Final (1999-2000)		
4. TITLE AND SUBTITLE Spatial coherence and crest-length statistics of waves		5. FUNDING NUMBERS N00014-99-11070		
6. AUTHOR(S) Steve Elgar & T.H.C. Herbers				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Washington State University Woods Hole Oceanographic Institution		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 North Quincy Arlington, VA 22217		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12. DISTRIBUTION / AVAILABILITY STATEMENT		12. DISTRIBUTION CODE Approved for public release		
13. ABSTRACT (Maximum 200 words) The methodology to simulate realizations of a nonlinear sea surface with a specified frequency-directional spectrum and a specified bispectrum have been developed. The simulation algorithm accounts for interactions between waves traveling in different directions. The target wavenumber spectrum can be obtained from observations, the output of a wave prediction model, or theory. The target bispectrum can be obtained from observations or from second-order nonlinear theory given the wavenumber spectrum. The bispectrum estimated from the wavenumber spectrum from LIDAR observations and second-order theory compares well with the bispectrum estimated directly from the images of approximately 1 m high waves in 20-30 m water depth near the North Carolina coast. Third-order statistics of the sea surfaces obtained from numerical simulations using the observed wavenumber spectrum and either the observed or theoretical bispectrum compare well with the statistics of the observed sea surface, verifying both the simulation technique and second-order nonlinear theory.				
14. SUBJECT TERMS waves, wave groups, sea surface wave simulations, crest lengths		15. NUMBER OF PAGES 3		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT unclassified	20. LIMITATION OF ABSTRACT unlimited	